



# What Is Double Diabetes?

by ANGELA FINLAY

---

## Causes and Treatment of Double Diabetes

Double diabetes is precisely what it sounds like: symptoms of type 1 diabetes occurring alongside symptoms of type 2 diabetes, making for a very uncomfortable situation.

It may seem like a bizarre phenomenon, but it's logical that those who have type 1 diabetes could develop the risk factors for type 2 diabetes, and eventually succumb to the insulin resistance that defines it.

The danger of double diabetes is also twofold.

Symptoms can be more numerous, more severe, and more difficult to control, but treatment also becomes more intricate to address both sets of symptoms. Prevention is extremely important, but there are ways to manage double diabetes, and possibly reverse some damage.

## What Causes Double Diabetes?

In recent years, the amount of double diabetes cases has increased, and it doesn't look like the trend will reverse anytime soon.

Since double diabetes is a relatively new health phenomenon, there aren't any firm statistics on the rate of growth, but recent studies have found that over 30 percent of newly diagnosed adolescent diabetics have both type 1 and type 2 of the disease.

## Defining Characteristics

Double diabetes is essentially severe insulin resistance.

Not only is the pancreas unable to produce enough insulin to support normal metabolism, but the body also stops responding to the injections of insulin that artificially encourage the important metabolic processes.

This puts the diabetic in a precarious position, since their body can no longer pull the glucose from the bloodstream and into the cells (it needs insulin to do that), and blood glucose levels stay elevated.

## Risk Factors

Obesity is the first and foremost risk factor for double diabetes.

It occurs when type 1 diabetics develop a resistance to the insulin they take by injection – just as a person with type 2 diabetes develops a resistance to the insulin, their pancreas naturally produces.

---

---

Not surprisingly, being overweight can also mask a proper diabetes diagnosis from the very start, since obesity is typically linked to type 2 diabetes. In turn, double diabetes can go undetected until the real cause – type 1 diabetes – is unveiled.

### **How it Begins**

Experts suspect that double diabetes strikes type 1 diabetics who take insulin to control their condition but haven't adopted any other lifestyle changes to manage their health.

Insulin use causes weight gain, which means those with type 1 diabetes must pay special attention to increasing their activity level and eating a healthy, calorie-controlled diet.

Failing to make these positive changes can introduce a new, extremely complicated illness as the body loses merely the ability to use insulin properly.

### **Other Causes of Insulin Resistance**

The classic cause of insulin resistance – and subsequently double diabetes, as we've discussed, is obesity. However, there are other reasons why people go on to develop insulin resistance – and sometimes, these are temporary situations.

For example, in times of extreme stress, such as intense physical, psychological or emotional stress, our bodies can become much more insulin resistant. A woman's body becomes much more insulin resistant during pregnancy. There are also a myriad of medications that cause insulin resistance.

According to the *Diabetes Sisters*, "long-term use of these types of medication may be necessary as treatment strategies for other health problems besides diabetes, and unintentionally cause insulin resistance in T1s and increased insulin resistance in T2s. Thus for some people, DD is medication induced."

"These types of medications" include corticosteroids, which are used to treat inflammation, certain psychotropic medications, and anti-seizure medications.

### **Controlling Double Diabetes Symptoms**

Type 1 and type 2 diabetes are treated differently, which makes a combined treatment plan somewhat complicated.

The first step is to determine the precise issues at hand, and the healthcare professionals can carefully create a medication regimen to help each set of symptoms.

### **Accurate Diagnosis**

It may seem like an easy distinction, but diagnosing the specific type of diabetes can be difficult.

Unfortunately, there seems to be a genetic predisposition connecting both forms of diabetes, and studies have shown that a large proportion of young people diagnosed with diabetes have the specific antibodies involved with type 1 diabetes, but also continue to produce insulin (which occurs with type 2 diabetes).

### **Insulin Therapy Adjustments**

Once the dual diabetes diagnosis has been confirmed, patients will need to inject insulin and take an oral medication to control both types of metabolic malfunctions.

Often the patients we've seen already have the type 1 diagnosis, so they have been injecting basal, correction, and prandial (insulin for food coverage) insulin. However, if a "double" diabetes diagnosis is confirmed, insulin

---

therapy adjustments will need to be made, meaning that there need to be increased to the insulin doses.

Also, a medication called Metformin is often prescribed in conjunction with insulin therapy – this helps insulin get into the cells. However, this is considered an “off-label” use of this medication and not every clinician will prescribe it. Research continues to address better approaches to counteracting insulin resistance.

### **Lifestyle Changes**

It's not too late to take matters into your own hands. Since weight gain is the primary problem behind double diabetes, weight loss is a good avenue to fewer complications.

A healthy diet is crucial, but muscle-building exercise can go far in the fight for better insulin response: lower blood sugar levels, more efficient organ function, and weight loss are some of the biggest benefits that come with regular activity.

### **The Dangers of Double Diabetes**

According to researchers in *Diabetologia*, the term “double diabetes” originated in 1991. It resulted when clinicians started noticing that people with type 1 diabetes who had a family history of type 2 diabetes were more likely to carry extra weight, and subsequently have a harder time reaching glycemic goals – even when they increased their insulin doses.

Since that time, there have been several research studies on double diabetes. Not only has the research found that patients with double diabetes have a harder time reaching glycemic goals, but they also are at a higher risk for complications.

For example, the EURODIAB study evaluated 3250 people with type 1 diabetes from 16 different countries. The study found that women who had a personal family history of type 2 diabetes were at a higher risk of developing albuminuria (protein in the urine – a risk factor for nephropathy) than those without a family history of type 2 diabetes.

A cross-sectional study from the Pittsburg Epidemiology of Diabetes Complications (EDC) cohort evaluated 658 patients, 112 of who had a family history of type 2 diabetes and 119 of whom had some event with their heart. Those who had a family history of type 2 diabetes were found to be at an increased risk for heart disease.

In a study performed by researchers in *Diabetologia*, the researchers concluded that when someone with type 1 diabetes begins to gain a significant amount of weight, clear action should be taken *before* insulin resistance sets in.

The researchers' state, “early consideration should be given to adjusting that regimen in the context of diet and lifestyle in an effort to limit weight gain. It may also be useful to ascertain a more complete family history of type 2 diabetes in such patients ... Lower thresholds (BP and lipid-lowering) for primary prevention of CVD should also be explored, as should the role of metformin as an ‘insulin-sparing’ agent.”

### **The Bottom Line...**

It is also important to remember that first and foremost, you must take care of *you*.

It may be hard to look at the big picture when you feel fine right now, but in a year or two, or even many years down the line, if you don't take your disease seriously, you could be facing complications.

Working hard to control your blood glucose levels – utilizing diet, exercise, medications, insulin, and the proper resources, such as a dietitian and a diabetes educator – can all help you achieve your goals.